

POSTOPERATIVE INFORMATION

FRANK A. PETRIGLIANO, MD

CHIEF – DIVISION OF SPORTS MEDICINE
HEAD TEAM PHYSICIAN – LA KINGS
ORTHOPAEDIC SURGERY AND SPORTS MEDICINE
USC DEPARTMENT OF ORTHOPAEDIC SURGERY
323.442.5822 OFFICE
323.865.5480 FAX

Keck Medicine
of USC



PATIENT STICKER

POST-OPERATIVE INSTRUCTIONS – ARTHROSCOPIC SHOULDER SURGERY--ACROMIOPLASTY / AC RESECTION

WOUND CARE

- You may remove the Operative Dressing on Post-Op Day #2. THERE ARE WHITE TAPES DIRECTLY ON THE SKIN. THESE ARE STERI-STRIPS—THESE SHOULD REMAIN ON.
- Apply Band-Aids to the wounds. Change the Band-Aids daily until the wounds are completely dry. Please do not use Bacitracin or other ointments under the bandage.
- Use Ice or cold therapy device 30-40 minutes 3-4 times per day until swelling is down, then as needed for pain relief.
- You may shower on Post-Op Day #2. Please cover the wound with plastic wrap and secure it to your skin with tape. You may remove the sling for showering, but keep the arm across the chest. Gently pat the area dry. Do not soak the shoulder in water. Do not go swimming in the pool or ocean until cleared by Dr. Petrigliano.

EXERCISES

- Wear the sling at all times except when doing your exercises. You may remove the sling for showering, but keep the arm across the chest. When your shoulder begins to feel comfortable out of the sling, you may remove it (3-7 days).
- Please perform the exercises on the attached Home Exercise Program 2-3 times daily:
- Pendulums and shoulder range of motion
 - Elbow / Hand / Wrist Range of Motion Exercises
 - Grip Strengthening
 - Scapular Tightening

POSTOP

- You will be given two prescriptions for pain relief for you to use post-operatively:
 - NAPROSYN, an anti-inflammatory medication, is to be used twice a day with breakfast and dinner, for the first seven days after surgery.
**** IF YOU HAVE AN ULCER HISTORY OR DIFFICULTY USING ANTI-INFLAMMATORIES OR ASPIRIN >> DO NOT USE**
 - VICODIN or PERCOCET, a strong narcotic, to be used only on an “as needed” basis for pain in addition to the Naprosyn.
- If you have any adverse effects with the medications, please call our office.
- If you develop a Fever (101.5), Redness or Drainage from the surgical incision site, please call our office to arrange for an evaluation.

POSTOPERATIVE INFORMATION

FOLLOW-UP

- Please call the office to schedule a follow-up appointment for your wound check, 7-10 days post-operatively. Dr. Petrigliano will prescribe physical therapy at this time.

IF YOU HAVE ANY QUESTIONS, PLEASE FEEL FREE TO CALL OUR OFFICE.