## **POSTOPERATIVE INFORMATION**

## FRANK A. PETRIGLIANO, MD

CHIEF – DIVISON OF SPORTS MEDICINE HEAD TEAM PHYSICIAN – LA KINGS ORTHOPAEDIC SURGERY AND SPORTS MEDICINE USC DEPARTMENT OF ORTHOPAEDIC SURGERY

323.442.5822 OFFICE 323.865.5480 FAX





PATIENT STICKER

## POST-OPERATIVE INSTRUCTIONS – KNEE ARTHROSCOPY

١	٨	ı		ı	ı	٨	ı	١,	^	۸	R	E
١	"	•	$\overline{}$	л.	_	ı١	ш	_	_	н	П	E

		You may remove the Operative Dressing on Post-Op Day #2. THERE ARE WHITE TAPES DIRECTLY ON TH								
	SKIN. THESE ARE STERI-STRIPS—THESE SHOULD REMAIN ON.									
		Apply Band-Aids to the wounds. Please <u>do not use</u> Bacitracin or other ointments								
		under the bandage. An ACE wrap may be used to help control swelling. Do not wrap the ACE too tight.								
		Use Ice or the Cryocuff as often as possible for the first 3-4 days, then as needed for pain relief. Do not								
		wrap the Ace too thickly or the cold may not penetrate.								
		You may shower on Post-Op Day #2. Cover the incisions with plastic while you shower. Gently pat the								
		area dry if it gets wet. <b>Do not soak</b> the knee in water. <b>Do not go swimming</b> in the pool or ocean until you								
		are cleared by Dr. Petrigliano to do so.								
		Crutches are provided for comfort. It is safe to put weight on the leg. When you feel comfortable								
		walking without the crutches you may do so (on average 3-5 days).								
<b>EXERCISES</b>	•									
		Keep your leg elevated with a pillow under your calf, NOT under the knee								
		Please perform the exercises on the attached Home Exercise Program 2-3 times daily:								
		Towel Roll Under Heel								
		Isometric Quadriceps Strengthening								
		Straight Leg Raise								
		<ul> <li>Seated Active Assisted Range of Motion Exercises</li> </ul>								
POST-OP		· ·								
		Enclosed are two prescriptions for you to use post-operatively:								
		• NAPROSYN, an anti-inflammatory medication, is to be used twice a day with breakfast								
		and dinner, for the first ten days after surgery.								
		** IF YOU HAVE AN ULCER HISTORY OR DIFFICULTY USING ANTI-INFLAMMATORIES OF								
		ASPIRIN >> USE ONLY VICODIN								
		<ul> <li>VICODIN, a strong narcotic, is to be used for pain relief as needed.</li> </ul>								
		If you have any adverse effects with the medications, please call our office.								
	☐ If you develop a Fever (101.5), Redness or Drainage from the surgical incision site, please of									
		to arrange for an evaluation.								
FOLLOW-U	JP									
		Please call the office to schedule a follow-up appointment for your wound check, 7-10 days post-								
		operatively. Dr. Petrigliano will prescribe physical therapy at this time.								

IF YOU HAVE ANY QUESTIONS, PLEASE FEEL FREE TO CALL OUR OFFICE.