

# POSTOPERATIVE INFORMATION

## FRANK A. PETRIGLIANO, MD

ORTHOPAEDIC SURGERY AND SPORTS MEDICINE  
UCLA DEPARTMENT OF ORTHOPAEDIC SURGERY  
310.319.1234 APPT  
310.825.2126 OFFICE  
310.825.1311 FAX  
CA License: A90515



PATIENT STICKER

### POST-OPERATIVE INSTRUCTIONS ULNAR COLLATERAL LIGAMENT RECONSTRUCTION

1. A plaster splint will be placed on your arm after surgery. You will not be able to bend or straighten your arm. However, gently make a loose fist and spread your fingers apart to keep the circulation moving in your arm.
2. Pain medication (PERCOCET) and an anti-inflammatory (NAPROSYN) are prescribed for you following surgery. Use the pain medication when needed. Use the anti-inflammatory as prescribed as it will help keep the swelling to a minimum and will help control pain.
3. You may shower. Cover the arm with a plastic bag and secure it around your upper arm with tape.
4. Your fingers may become swollen. Moving them frequently and elevating your arm above your heart will help minimize this.
5. You may notice SLIGHT numbness/tingling in your fingers. This is normal for the first 12-24 hours. If this persists or worsens, contact the office. You may be directed to loosen the ace bandage on your arm.
6. You may use your hand to write. You may NOT lift or push anything. Because the elbow is nearly straight, it may be difficult to use it to eat, etc.
7. Because the splint is heavy, your shoulder may become sore. Shrug your shoulders frequently to help minimize this.
8. Please call the office to schedule a follow-up appointment for wound check and application of brace, 10 days post-operatively.

**IF YOU HAVE ANY QUESTIONS, PLEASE FEEL FREE TO CALL OUR OFFICE.**